

REPORT DOCUMENTATION PAGEForm Approved
OMB NO. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comment regarding this burden estimates or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 6/30/2000	3. REPORT TYPE AND DATES COVERED Final Report 7/1/98-6/30/00	
4. TITLE AND SUBTITLE Intersubband Lasers at Infrared and Terahertz Frequencies			5. FUNDING NUMBERS DAAG55-98-1-0434	
6. AUTHOR(S) G. I. Haddad				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) G. I. Haddad Electrical Engineering and Computer Science Department University of Michigan Ann Arbor, MI 48109-2122			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAMES(S) AND ADDRESS(ES) U.S. Army Research Office P.O. BOX 12211 Research Triangle Park, NC 27709-2211			10. SPONSORING / MONITORING AGENCY REPORT NUMBER ARO 38858.4-EL	
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) This research project was aimed at an investigation of far-infrared (FIR) and terahertz (THz) sources using intersubband transitions in step quantum well structures. Theoretical studies were carried out in order to determine the optical gain spectra. The intersubband gain was formulated as a function of the dipole matrix elements, the lineshape function, and the electron distribution functions in the lasing subbands. The dependence of the spectral gain on optical phonon scattering was investigated. Non-Lorentzian lineshape functions were obtained and it was found that the magnitude and overall shape of the spectral gain are significantly modified by the intra- and inter-subband phonon scattering rates. Effects of the confined phonon modes were evaluated and it was found that they have a significant effect on the devices considered.				
14. SUBJECT TERMS Intersubband Lasers; Confined Phonons; Quantum Wells			15. NUMBER OF PAGES	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OR REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

Final Report

**Intersubband Lasers at Infrared
and Terahertz Frequencies**

Submitted to

Army Research Office
P. O. Box 12211
Research Triangle Park, NC 27709-2211

Attention: Dr. Dwight Woolard

Submitted by

Professor George I. Haddad
Solid State Electronics Laboratory
Department of Electrical Engineering and Computer Science
The University of Michigan
Ann Arbor, MI 48109-2122

Telephone: 734-764-7139
Fax: 734-763-9324
Email: gih@eecs.umich.edu

July 1, 2000

20001122 146

Intersubband Lasers at Infrared and Terahertz Frequencies

July 1, 1998-June 30, 2000

1. Abstract:

This research project was concerned with a theoretical investigation of the properties of intersubband lasers for Far Infrared (FIR) and Terahertz (THz) frequencies. Theoretical models were developed to determine the optical gain spectra in various types of intersubband lasers including Quantum Cascade (QCL) and Step Quantum Well Lasers (SQL). The dependence of device properties on various material parameters and the effects of confined phonon modes were investigated. The results obtained from these studies will be very important in the design of lasers based on intersubband transitions.

2. Scientific Personnel:

Faculty Supervisor: Prof. G. I. Haddad
Research Staff: Dr. J. P. Sun
Graduate Students: None

3. Specific Aim:

The specific aim of this project was to develop improved models for predicting spectral gain in intersubband lasers. This included a determination of phonon relaxation rates which are very important in determining the properties of such lasers. In particular, special consideration was given to the effects of confined phonon modes in quantum wells and their importance in determining basic device design and properties.

4. Significant Accomplishments:

- Modeling capability of various laser physical parameters was developed to establish a design capability for intersubband laser structures. The calculations based on these models included the electron states in intersubband laser structures, intersubband relaxation rates, tunneling injection and escape times, carrier population inversion, transition matrix elements, oscillator strength, lineshape functions, optical gain spectra, threshold current density, and output power. This is significant in designing optimum device schemes for FIR and THz lasers.
- We have also studied effects of rough heterointerfaces on the shape of the THz absorption peak in quantum well structures. The equation for intersubband polarization was considered in the resonant approximation, taking into account the depolarization shift. The lineshape of the intersubband absorption peak was formulated and calculated for the case with long range variations of heterointerfaces.

- The intersubband optical gain was formulated and calculated for two step quantum well laser structures under investigation. This is an important step in the design of optimum device schemes for FIR and THz lasers since the gain determines the laser threshold characteristics. It is calculated as a function of several critical device design parameters, including electron population inversion, distribution functions in the subbands, the lineshape function, temperature, and photon frequency.
- We have calculated non-Lorentzian lineshape functions for these laser structures depending on various intra- and inter-subband phonon scattering mechanisms. This is significant progress as compared to the usual adoption of the Lorentzian lineshapes which may lead to either an over-estimate or under-estimate of the optical gain. Moreover, the confined phonon effects, especially effects of the interface optical phonon scattering on the gain have been included, which can greatly modify the overall shape and magnitude of the gain. This will aid in the design of FIR and THz lasers.
- Our modeling capability has enabled us to study and conceive alternative device schemes for FIR and THz sources. We have investigated Quantum Cascade Lasers (QCL) and Step Quantum Well Lasers (SQL) which were conceived by us and may result in enhanced performance as compared to QC Lasers.
- The results of this investigation are given in detail in the following publications which are attached as part of this report.

5. Publications:

H. B. Teng, J. P. Sun, G. Haddad, M. Stroscio, S. Yu and K. Kim, "Phonon Assisted Intersubband Transitions in Step Quantum Well Structures," *J. Appl. Phys.*, **84**: 2155-2164, 1998.

M. A. Stroscio, "Quantum Cascade Lasers and Quantum Well Intersubband Lasers," a book chapter in *Advances in Semiconductor Lasers and Applications to Optoelectronics*, M. A. Stroscio and M. Dutta, eds., to be published,

F. T. Vasco, J. P. Sun, G. I. Haddad and V. V. Mitin, "Inhomogeneous Broadening of Intersubband Transitions Due to Nonscreening Roughness of Heterointerfaces," to appear in *J. Appl. Phys.*

6. Honors and Awards:

Prof. Haddad received the Third Millennium Medal from the IEEE-Microwave Theory and Techniques Society for his significant contributions to the profession.